

What is **Non-Small-Cell Lung Cancer?**

Let us answer some
of your questions.

Non-small-cell lung cancer (NSCLC)

An ESMO guide for patients

Patient information based on ESMO Clinical Practice Guidelines

This guide has been prepared to help you, as well as your friends, family and caregivers, better understand the nature of non-small-cell lung cancer (NSCLC) and the treatments that are available. It includes information on the different subtypes of NSCLC, the causes of the disease and how it is diagnosed, and up-to-date guidance on the types of treatments that may be available and any possible side effects.

The medical information described in this document is based on the ESMO Clinical Practice Guidelines (CPG) for NSCLC, which are designed to help medical oncologists with the diagnosis and management of early stage, locally advanced and metastatic NSCLC. All ESMO CPGs are prepared and reviewed by leading experts using evidence gained from the latest clinical trials, research and expert opinion.

The information included in this guide is not intended as a replacement for your doctor's advice. Your doctor knows your full medical history and will help guide you regarding the best treatment for you.

This guide has been developed and reviewed by:

Representatives of the European Society for Medical Oncology (ESMO):

Silvia Novello; Solange Peters; Raffaele Califano; Jean-Yves Douillard; Francesca Longo; Claire Bramley; and Svetlana Jezdic

Representatives of the European Oncology Nursing Society (EONS):

Anita Margulies; Roisin Lawless

Patient advocate from the Lung Cancer Europe (LuCE) and Women Against Lung Cancer in Europe (WALCE): Stefania Vallone

2	An ESMO guide for patients
4	Lung cancer: A summary of key information
7	Anatomy of the lungs
8	What is lung cancer?
9	How common is NSCLC?
11	What causes NSCLC?
13	How is NSCLC diagnosed?
16	How will my treatment be determined?
20	What are the treatment options for NSCLC?
32	What are the possible side effects of treatment?
42	What happens after my treatment has finished?
44	Support groups
45	References
47	Glossary

Lung cancer: A summary of key information

Introduction to lung cancer

- Lung cancer arises from cells in the lung that have grown abnormally and multiplied to form a lump or **tumour**.
- Non-small-cell lung cancer (NSCLC) is a type of lung cancer, which is differentiated from small-cell lung cancer (SCLC) because of the way the **tumour** cells look under a microscope. The three main types of NSCLC are **squamous cell carcinoma**, **adenocarcinoma** and **large cell (undifferentiated) carcinoma** of the lung. They are diagnosed in the same way but may be treated differently.
- Lung cancer is the fourth most common cancer in Europe; NSCLC represents 85-90% of all lung cancers. Smoking is the biggest risk factor for the development of lung cancer.
- In Europe, there has been a decrease in lung cancer mortality among men, while it is increasing in women – this reflects a difference in smoking trends between the sexes.

Diagnosis of NSCLC

- Lung cancer may be suspected if a person has symptoms such as persistent cough or chest infection, breathlessness, hoarseness, chest pain or coughing up blood. Other symptoms may be fever, appetite loss, unexplained weight loss and **fatigue**.
- Following a clinical examination, your doctor will arrange for an **x-ray** and/or **computed tomography** (CT) scan (or might use other technologies, such as positron emission tomography [PET] CT scan or magnetic resonance imaging [MRI]) to evaluate the position and extent of the cancer. Examination of a **biopsy** (cells or tissue taken from the **tumour**) will confirm a diagnosis of NSCLC.

Treatment options for NSCLC

- Types of treatment include:
 - Surgery
 - **Radiotherapy** – the use of measured doses of radiation to damage cancer cells and stop them growing.
 - **Chemotherapy** – the use of anti-cancer drugs to destroy cancer cells. **Chemotherapy** can be given alone or with other treatments.
 - **Targeted therapy** – newer drugs that work by blocking the signals that tell cancer cells to grow.
 - **Immunotherapy** – a type of treatment designed to boost the body's natural defenses in order to fight cancer.
- Combinations of different treatment types are frequently offered based on the stage and type of NSCLC and on the patient's condition and **comorbidities** (additional diseases or disorders experienced at the same time). **Adjuvant** treatment – the use of anticancer drugs after or in combination with another form of cancer treatment – may be used in some patients.

- Cancer is 'staged' according to **tumour** size, involvement of **regional lymph nodes** and whether it has spread outside the lung to other parts of the body. This information is used to help decide the best treatment.
- **Early-stage (stage I-II) NSCLC:**
 - Surgery is the main treatment for **early-stage** NSCLC.
 - **Chemotherapy** may be given after surgery (**adjuvant chemotherapy**) in patients with stage II NSCLC.
 - **Radiotherapy** (either **stereotactic ablative radiotherapy** [SABR] or **conventional radiotherapy**) is an alternative to surgery in patients who are unable or unwilling to have surgery.
 - **Adjuvant radiotherapy** may be given when it has not been possible to completely remove the **tumour** during surgery.
- **Locally advanced (stage III) NSCLC:**
 - Treatment is likely to involve different types of therapy (**multimodal therapy**).
 - If it is possible to remove the **tumour** (i.e. the **tumour** is **resectable**), treatment options include:
 - ~ **Induction therapy** (initial treatment[s] given to shrink the **tumour** before a second planned treatment) consisting of **chemotherapy** with or without **radiotherapy**, followed by surgery.
 - ~ Surgery followed by **adjuvant chemotherapy** and/or **radiotherapy**.
 - ~ **Chemoradiotherapy** (i.e. **chemotherapy** and **radiotherapy** given at the same time).
 - The type of treatment – and sometimes the sequence of treatments – offered to patients with **resectable** stage III NSCLC will depend on the extent and complexity of the surgery required to remove the **tumour**.
 - In **unresectable** stage III NSCLC, **chemoradiotherapy** is the preferred treatment. Alternatively, **chemotherapy** and **radiotherapy** can be given **sequentially** (i.e. one after the other) in patients unable to tolerate **concurrent** treatment.
- **Metastatic (stage IV) NSCLC:**
 - NSCLC is referred to as **metastatic** or stage IV disease when it has spread beyond the lung which was initially affected.
 - It is rarely possible to remove **metastatic** NSCLC with surgery or to treat it radically with **radiotherapy**.
 - Intravenous **chemotherapy** with a two-drug combination (with or without the addition of the **targeted therapy** called **bevacizumab**) is the main treatment for patients with **metastatic** NSCLC. New options using **first-line immunotherapy** are under evaluation and **pembrolizumab** has recently been approved for use in this setting. Thus, **immunotherapy** is likely to replace or complement **first-line chemotherapy** in selected patients in the next few years.
 - The choice of drugs used will largely depend on the general health of the patient and the **histological subtype** of the **tumour**.

- Patients whose **tumours** contain specific **mutations** (alterations) to the **epidermal growth factor receptor (EGFR)** or **anaplastic lymphoma kinase (ALK)** genes (determined by molecular testing using a **tumour biopsy**) are best treated with oral **targeted therapies** given continuously.
- After 4-6 cycles of **doublet chemotherapy** (i.e. two **chemotherapy** drugs given together), **maintenance treatment** (treatment to help keep the cancer from coming back) with a **chemotherapy** drug called **pemetrexed**, with or without a **targeted therapy** called **bevacizumab**, may be given to patients in good general health. The **targeted therapy erlotinib** may be offered as **maintenance treatment** in patients whose **tumours** have **EGFR mutations**.
- Should the cancer come back (**relapse** or recurrence), **second-** and **third-line** treatments may be offered. Suitable **second-** and **third-line** treatments depend on which **first-line** treatment has been received and on the general health of the patient. Treatment options include: **Chemotherapy** (**pemetrexed** or **docetaxel**), **immunotherapy** (**nivolumab** or **pembrolizumab**), **antiangiogenic therapy** (**nintedanib** or **ramucirumab**) in combination with **docetaxel**, and **targeted therapies** (**afatinib**, **gefitinib**, **erlotinib**, **crizotinib**, **ceritinib**, **alectinib** or **osimertinib**).
 - ~ The activity of **immunotherapy** is influenced by the amount of **programmed death-ligand 1 (PD-L1)** protein in the **tumour** (determined by molecular testing using a **tumour biopsy**). **Pembrolizumab** – but not **nivolumab** – can only be prescribed in **tumours** that are **PD-L1** positive. Other **immunotherapy** drugs are under evaluation in **clinical trials**.
 - ~ Patients whose **tumours** have **EGFR mutations** who have received **first-line** treatment with **erlotinib**, **gefitinib** or **afatinib**, and who have a confirmed **EGFR T790M mutation**, may be subsequently treated with **osimertinib**.
 - ~ Patients whose **tumours** have **ALK** rearrangements and who have received **first-line** treatment with **crizotinib** may be treated with **second-line ceritinib** or **alectinib**. Other **ALK** inhibitors are under evaluation in **clinical trials**.


Follow-up after treatment

- Patients who have completed treatment for stage I-III NSCLC are followed-up with clinical and **radiological examinations** every 3-6 months for the first 2-3 years and annually after that.
- Patients who have completed treatment for **metastatic** disease are followed up with **radiological examinations** every 6-12 weeks (depending on their suitability for further treatment) so that **second-line** therapy can be started if needed.

Anatomy of the lungs

The lungs form part of our respiratory (breathing) system, which includes:

- Nose and mouth.
- **Trachea** (windpipe).
- **Bronchi** (tubes that go to each lung).
- Lungs.



*Anatomy of the respiratory system, showing the **trachea**, **bronchi**, and lungs. As we breathe in, air passes from our nose or mouth, through the **trachea**, **bronchi** and **bronchioles**, before it reaches tiny air sacs called **alveoli** – this is where oxygen from the air passes into the bloodstream (see inset image).*

What is lung cancer?

Lung cancers typically start in the cells that line the **bronchi** and parts of the lung such as the **bronchioles** or **alveoli**. There are two main types of **primary lung cancer**:

- **Small-cell lung cancer (SCLC)**: This type gets its name from the small size of the cells that it is composed of when viewed under a microscope.
- **Non-small-cell lung cancer (NSCLC)**: This is the more common type of lung cancer, and accounts for 85–90% of all lung cancers (*Novello et al., 2016*).
 - This guide will focus exclusively on NSCLC.

What subtypes of NSCLC are there?

The three main **histological subtypes** of NSCLC are:

- **Adenocarcinoma**: About 40% of all lung cancers are **adenocarcinomas**. These **tumours** start in mucus-producing cells that line the airways.
- **Squamous cell carcinoma (SCC)**: About 25–30% of all lung cancers are **SCC**. This type of cancer develops in cells that line the airways and is usually caused by smoking.
- **Large cell (undifferentiated) carcinoma**: This type makes up around 10–15% of all lung cancers. It gets its name from the way that the cancer cells look when they are examined under a microscope.

What are the symptoms?

The most common symptoms of lung cancer, including NSCLC, are:

- Persistent cough.
- Coughing blood.
- Chest infection that won't go away or keeps coming back.
- Chest or shoulder pain that won't go away.
- Difficulty breathing/shortness of breath.
- Hoarseness or lowering of the voice.
- Wheezing.

Other, non-specific symptoms, may include:

- Fever
- Loss of appetite.
- Unexplained weight loss.
- Feeling extremely tired.

You should see your doctor if you experience any of these symptoms. However, it is important to remember that these symptoms are common in people who do not have lung cancer; they may also be caused by other conditions.

How common is NSCLC?

Lung cancer represents the fourth most common cancer in Europe

In 2012, there were more than 410,000 new cases of lung cancer diagnosed in Europe (12% of the total number of new cancer cases) (*Ferlay et al., 2013*):

- 291,000 new cases in men.
- 119,000 new cases in women.

Lung cancer is the second most common cancer in men (after prostate cancer) and the third most common in women (after breast and colorectal cancer) (*Ferlay et al., 2013*). Incidence rates of lung cancer are higher in more developed countries than in less developed countries; these variations largely reflect the differences in the stage and degree of the tobacco epidemic (*Torre et al., 2015*).


In Europe, there has been a decrease in lung cancer mortality among men, while it is increasing in women – this reflects the difference in smoking prevalence trends between the sexes (*Malvezzi et al., 2016, Novello et al., 2016*).

The majority of cases of lung cancer are diagnosed in patients aged 65 years and over, and the median age at diagnosis is 70 years.

NSCLC is the most common type of lung cancer, representing 85-90% of all lung cancers

Non-small-cell lung cancer

Estimated lung cancer incidence rates by sex and world area in 2012 (Torre et al., 2015).



What causes NSCLC?

Smoking is the biggest risk factor for developing lung cancer. However, there are other risk factors that can also increase the chances of developing lung cancer. It is important to remember that having a risk factor increases the risk of cancer developing but it does not mean that you will definitely get cancer. Likewise, not having a risk factor does not mean that you definitely won't get cancer.

Smoking is the biggest risk factor for lung cancer

Smoking

Tobacco smoking is the leading cause of lung cancer. In Europe, it is responsible for 90% of cases in men and 80% of cases in women (*Novello et al., 2016*). The number of years that a person has been a smoker is more important than the number of cigarettes smoked per day; therefore, giving up smoking at any age can reduce the risk of developing lung cancer more than cutting down on the number of cigarettes smoked per day.




Passive smoking

Passive smoking, also referred to as 'second-hand smoke' or 'environmental tobacco smoke', increases the risk of developing NSCLC but to a lesser extent than if you are a smoker.



Radon

Radon is a **radioactive** gas that is produced during the breakdown of naturally-occurring **uranium** in soil and rocks, particularly granite. It can pass through from the ground into homes and buildings. Exposure to excessive levels of radon is thought to be a significant causative factor in patients with lung cancer who have never smoked. This may be particularly relevant for underground miners who are usually exposed to high levels of radon if the mines in which they work are in a particular geographical region.



Genetic susceptibility

It is thought that some people may be more likely to develop lung cancer based on their genetic makeup (*Bailey-Wilson et al., 2004*). Having a family history of lung cancer, or other types of cancer, increases the risk of developing lung cancer to some degree. In people who are genetically predisposed to lung cancer, smoking further increases the risk.



Household and environmental pollutants

Other factors described as risk factors for the development of NSCLC include exposure to **asbestos** and **arsenic**. There is evidence that lung cancer rates are higher in cities than in rural areas, although factors other than outdoor air pollution may be responsible for this pattern. It has also been suggested that indoor air pollution from use of coal-fuelled stoves may be a factor in some countries (*Novello et al., 2016*). For example, in China there is an increased rate of lung cancer in women, despite the fact that a lower proportion of women are smokers in China compared with some European countries.



At the present time, there is no clear evidence that screening for NSCLC should be a routine procedure in people who are at a higher risk of developing the disease based on the above risk factors.

How is NSCLC diagnosed?

Most patients with NSCLC are diagnosed after seeing their doctor to report symptoms such as a persistent cough, a chest infection that won't go away, **dyspnoea**, wheezing, coughing blood, chest or shoulder pain that won't go away, hoarseness or lowering of the voice, unexplained weight loss, loss of appetite or extreme **fatigue**.

A diagnosis of lung cancer is based on the results of the following examinations and tests:

Clinical examination

Your doctor will carry out a clinical examination. He/she will examine your chest and check for the **lymph nodes** in your neck. If there is a suspicion of lung cancer, he/she may arrange for a chest **x-ray**, or possibly a **CT** scan, and refer you to a specialist for further testing.



Imaging

Imaging is used to confirm a suspected diagnosis of lung cancer, and to investigate how far the cancer has progressed

Different imaging techniques include:

- **Chest x-ray:** A chest **x-ray** will enable the specialist to check your lungs for anything that looks abnormal. This is usually the first test that is carried out, based on your symptoms and the clinical examination.
- **CT scan of chest and upper abdomen:** A series of images are taken, which build up a three-dimensional picture of the inside of your body. This allows the specialist to gather more information about the cancer such as the exact location of the **tumour** in your lungs, whether nearby **lymph nodes** are affected, and whether the cancer has spread to other areas of the lungs and/or parts of your body. It is a painless procedure and usually takes about 10-30 minutes.
- **CT scan or magnetic resonance imaging (MRI) scan of the brain:** This test allows doctors to rule out or confirm whether the cancer has spread to your brain. An **MRI** scan uses powerful magnetism to build up detailed images. You may be given an injection of dye into a vein in your arm to help the images show up more clearly. The scan won't hurt but may be slightly uncomfortable as you will need to lie still inside the scanning tube for about 30 minutes. You will be able to hear and speak to the person doing the scan.



- **Positron emission tomography (PET)/CT scan:** A combination of a **CT** scan and a **PET** scan. **PET** uses low-dose radiation to measure the activity of cells in different parts of the body, so a **PET/CT** scan gives more detailed information about the part of the body being scanned. A mildly **radioactive** drug will be injected into a vein in the back of your hand or arm and then you will need to rest for about an hour while it spreads throughout your body. The scan itself will take 30-60 minutes and, although you will need to lie still, you will be able to speak to the person operating the scanner. A **PET/CT** scan is often carried out to detect whether the cancer has spread to the bones.


Histopathology

Examination of a biopsy is recommended for all patients with NSCLC as it helps to determine the best treatment approach

Histopathology is the study of diseased cells and tissues using a microscope; a **biopsy** of the **tumour** allows a sample of cells to be closely examined. Examination of a **biopsy** is recommended for all patients as it is used to confirm a diagnosis of NSCLC, to identify the **histological subtype** of NSCLC, and to identify any abnormal proteins within the **tumour** cells that could help to determine the best treatment for you (Novello *et al.*, 2016).

Techniques for obtaining a **biopsy** include:

- **Bronchoscopy:** A doctor or specially-trained nurse examines the insides of the airways and lungs using a tube called a **bronchoscope**. It is carried out under local anaesthetic. During a **bronchoscopy**, the doctor or nurse will take samples of cells (**biopsies**) from the airways or lungs.
- **CT-guided needle lung biopsy:** If a **biopsy** is difficult to obtain with a **bronchoscopy**, your doctor may choose to obtain a **biopsy** during a **CT** scan. In this procedure, you will have a **local anaesthetic** to numb the area. A thin needle is then inserted through your skin into your lung so that the doctor can remove a sample of cells from the **tumour**. This should only take a few minutes.
- **Endobronchial ultrasound-guided sampling (EBUS):** This technique is used to confirm whether the cancer has spread to nearby **lymph nodes**, after **radiological examinations** have suggested that this might be the case. A **bronchoscope**, containing a small **ultrasound**



probe, is passed through the **trachea** to see whether any nearby **lymph nodes** are larger than normal. The doctor can pass a needle along the **bronchoscope** to take **biopsies** from the **tumour** or the **lymph nodes**. This test can be uncomfortable but shouldn't be painful. It takes less than an hour and you should be able to go home the same day after it is finished.

- **Oesophageal ultrasound-guided sampling (EUS):** Similar to EBUS, this technique is used to confirm whether the cancer has spread to nearby **lymph nodes**, after **radiological examinations** have suggested that this might be the case. However, unlike EBUS, the **ultrasound** probe is inserted through the **oesophagus**.
- **Mediastinoscopy:** This procedure is more invasive than EBUS/EUS but is recommended as an extra test if EBUS/EUS does not confirm that the cancer has spread to nearby **lymph nodes** or if the **lymph nodes** requiring investigation cannot be reached by EBUS. A mediastinoscopy is carried out under **general anaesthetic** and requires a short stay in hospital. A small cut is made in the skin at the front of the base of your neck and a tube passed through the cut into your chest. A light and a camera attached to the tube allow the doctor to closely look at the middle of your chest – the mediastinum – for any abnormal **lymph nodes**, as these are the first areas that the cancer may spread to. Samples of tissue and **lymph nodes** can be taken for further examination.

Ask your doctor for details if you have any questions about these procedures

Cyto(patho)logy

Whereas histopathology is the laboratory examination of tissue or cells, cytology (or cytopathology) is the examination of cancerous cells spontaneously detached from the **tumour**. Common methods for obtaining samples for cytological examination include:

- **Bronchoscopy:** Bronchial washings (in which a mild salt solution is washed over the surface of the airways) and the collection of secretions can be carried out during a **bronchoscopy** to look for the presence of cancerous cells.
- **Thoracentesis/pleural drainage:** Pleural effusion is an abnormal collection of fluid between the thin layers of tissue (**pleura**) that line the lung and the wall of the chest cavity. This fluid can be taken from the pleural cavity by thoracentesis or pleural drainage and examined in the laboratory for the presence of cancerous cells.
- **Pericardiocentesis/pericardial drainage:** Pericardial effusion is an abnormal collection of fluid between the heart and the sac that surrounds the heart (**pericardium**). This fluid can be taken from the pericardial cavity by pericardiocentesis or pericardial drainage and examined in the laboratory for the presence of cancerous cells. These techniques are carried out in the hospital, usually with the aid of **ultrasound** to help position the needle. You will be given a **local anaesthetic** and monitored closely for any complications afterwards.

Because of the location of your lungs in your body, obtaining samples of cells/tissue can be difficult and it may be necessary to repeat some of these tests if results are found to be inconclusive.

How will my treatment be determined?

After a diagnosis is confirmed, your cancer specialist will look at a number of factors to help plan your treatment. This includes information about yourself and about the cancer.

Patient-related factors

- Your age.
- Your general health.
- Your medical history.
- Your smoking history.
- Results of blood tests and scans.

Cancer-related factors

Treatment also depends on the type of lung cancer that you have (histopathology or cytopathology results), where it is in the lung (its location) and whether it has spread to other parts of the body (imaging results).

Staging

It is important for your doctor to know the stage of the cancer so that he/she can determine the best treatment approach

Staging of the cancer is used to describe its size and position and whether it has spread from where it started. Cancer is staged using a number/letter system – described as stages IA-IV. Generally, the lower the stage the better the **prognosis**. Staging considers:

- How big the cancer is (**tumour** size).
- Whether it has spread into the **lymph nodes**.
- Whether it has metastasised (spread) to other areas within the lungs or to other parts of the body.

Staging is usually carried out twice: After clinical and **radiological examinations**; and after surgery, in the case of surgically resected **tumours**.

The different stages of NSCLC are described in the table below.

STAGE IA	<ul style="list-style-type: none"> The tumour is no larger than 3 cm, is still inside the lung and has not spread to any of the nearby lymph nodes 	Early-stage NSCLC
STAGE IB	<ul style="list-style-type: none"> The tumour is 3–5 cm in size, is still inside the lung and has not spread to any of the nearby lymph nodes 	
STAGE IIA	<ul style="list-style-type: none"> The tumour is 5–7 cm in size, is still inside the lung and has not spread to any of the nearby lymph nodes; or The tumour is no larger than 5 cm, has spread to nearby lymph nodes but is not in any other part of the body 	
STAGE IIB	<ul style="list-style-type: none"> The tumour is 5–7 cm in size, has spread to nearby lymph nodes but is not in any other part of the body; or The tumour is larger than 7 cm or there is more than one tumour in the same lobe; it has not spread to nearby lymph nodes but may invade other parts of the lung, the airway or the surrounding areas just outside the lung, e.g. the diaphragm 	
STAGE IIIA	<ul style="list-style-type: none"> The tumour is no larger than 7 cm, has spread to nearby lymph nodes but is not in any other part of the body; or The tumour is larger than 7 cm or there is more than one tumour in the same lobe; it has spread to nearby lymph nodes and may invade other parts of the lung, the airway or the surrounding areas just outside the lung, e.g. the diaphragm; or The tumour is of any size and invades tissues and structures further away from the lung, such as the heart, windpipe or oesophagus, but it has not spread to other parts of the body; or there is more than one tumour in different lobes of the same lung. The cancer may or may not have spread to nearby lymph nodes 	Locally advanced NSCLC
STAGE IIIB	<ul style="list-style-type: none"> The tumour is of any size and invades tissues and structures further away from the lung, such as the heart, windpipe or oesophagus, but it has not spread to other parts of the body; or there is more than one tumour in different lobes of the same lung; the cancer has also spread to nearby lymph nodes; or The tumour is of any size and may or may not invade tissues and structures further away from the lung, such as the heart, windpipe or oesophagus; or there is more than one tumour in different lobes of the same lung; the cancer has spread to more lymph nodes, but not to other parts of the body 	
STAGE IV	<ul style="list-style-type: none"> The tumour is of any size and may or may not have spread to the lymph nodes. The cancer is in both lungs, has spread to another part of the body (e.g. the liver, adrenal glands, brain or bones) or it has caused a collection of fluid around the lung or heart that contains cancer cells. Metastases are present either at diagnosis (in nearly 40% of patients) or they are discovered during follow-up tests of previously treated NSCLC 	Metastatic NSCLC

AJCC/UICC system 7th edition – abridged version (Novello et al., 2016)

An 8th edition of the AJCC/UICC staging system has been published, which contains detailed descriptions of **tumour** size and classification of **metastatic** disease according to whether it is **oligometastatic**, thoracic (in the chest) or widespread (AJCC, 2016).

AJCC, American Joint Committee on Cancer; NSCLC, non-small-cell lung cancer; UICC, Union for International Cancer Control

Type of NSCLC

Biopsy results

Your **biopsy** will be examined in the laboratory to determine:

- The **histological subtype** (**SCC**, **adenocarcinoma** or **large cell carcinoma**).
- **Grade**.
- **Tumour** biology.

Histological subtype

The **histological subtype** of the **tumour** can influence the type of treatment you will receive. For example, non-squamous cancers may benefit from certain anti-cancer therapies that have been shown to be effective only in patients with this **histological subtype**.

Grade

Grade is based on how different **tumour** cells look from normal lung cells, and on how quickly they grow. The **grade** will be a value between one and three and reflects the aggressiveness of **tumour** cells; the higher the **grade**, the more aggressive the **tumour**.

Biological testing of the tumour

Tissue specimens from **metastatic** NSCLC belonging to the non-squamous subtype should be tested for the presence of specific **mutations** in the **EGFR** gene. Even though such **mutations** are rare (approximately 10–12% in Caucasians with **adenocarcinoma**), the detection of an **EGFR** gene **mutation** has important prognostic and therapeutic implications in patients with **metastatic** NSCLC. **EGFR** testing is not recommended in patients with a diagnosis of **SCC**, except in never/former light smokers (<15 pack years) (Novello *et al.*, 2016).

Routine testing for rearrangement in the **ALK** gene is now standard of care and should be carried out, if possible, in parallel with **EGFR mutation** analysis. **ALK** rearrangement is more frequent in people who have never smoked, those with the **adenocarcinoma** subtype (5%) and in younger patients (aged <50 years old). Detecting **ALK** rearrangements has important therapeutic implications for patients with **metastatic** NSCLC due to the existence of drugs targeting **ALK** (e.g. **crizotinib**, **ceritinib** and **alectinib**) (Novello *et al.*, 2016).

Programmed death-ligand 1 (PD-L1): This is a cellular protein thought to be involved in helping the **tumour** to evade detection by the body's immune system. The amount of **PD-L1** present in a **tumour** may influence the decision to treat the cancer with anti-PD-L1 **immunotherapy**.

Who is involved in planning my treatment?

In most hospitals, a team of specialists will plan the treatment they feel is best for your individual situation. This **multidisciplinary team** of medical professionals may include:

- A surgeon.
- A medical **oncologist** (a doctor who specialises in the medical management of cancer).
- A radiation **oncologist**.
- A chest physician.
- A **nurse specialist**.
- A **radiologist** (or radiographer) who has been involved in the assessment of any **x-rays** and scans.
- A **pathologist** who has been involved in the analysis of your **tumour biopsy**.



Other services that may be offered include: a dietician, a social worker, a community care nurse, a physiotherapist, a clinical psychologist and a palliative care service (who can assist with pain management). After consultation with the **multidisciplinary team**, your doctor, possibly with other members of the care team, will talk to you about the best treatment plan for your situation (*Novello et al., 2016*). They will explain the benefits and potential drawbacks of different treatments.

What are the treatment options for NSCLC?

Aims of treatment

In **early-stage** NSCLC, when the cancer is confined to the lung and therefore considered to be curable, the main treatment is surgical resection (*Vansteenkiste et al., 2013; Vansteenkiste et al., 2014*). For **locally advanced** NSCLC, **multimodal therapy** is usually adopted to help shrink or in some cases completely remove the cancer (*Eberhardt et al., 2015*). For **metastatic** NSCLC, when the cancer has spread to other parts of the body, and cure is not an option, various **systemic treatments** may be used in an attempt to slow down **tumour** growth and improve symptoms and quality of life – this is called **supportive** or **palliative care** (*Novello et al., 2016*).

Overview of treatment types

Treatments for NSCLC include surgery, radiotherapy, chemotherapy and targeted therapies

The treatment you receive will depend on the stage and type of cancer, as well as your general health and treatment preferences, which will be discussed together with your doctor. You may have a combination of treatments. The main types of treatment are listed below:

- **Surgery** may be possible to remove NSCLC if it is diagnosed at an early stage. The type of operation that is offered will depend on the size and location of the cancer (*Vansteenkiste et al., 2013; Vansteenkiste et al., 2014*):
 - A **wedge** or **segment resection** is the removal of a very small amount of the lung; this is sometimes offered if the cancer is at a very early stage
 - A **lobectomy** is the removal of one of the **lobes** of the lung; it is the standard surgical treatment for NSCLC
 - A **pneumonectomy** is the total removal of one of the lungs; it is a more complex surgical **resection** than **lobectomy** or **wedge (segment) resection**.
- **Chemotherapy** works by disrupting the way that cancer cells grow and divide. However, these drugs can also affect normal cells. **Chemotherapy** can be given before or after surgery for NSCLC. Some people have **chemotherapy** at the same time as **radiotherapy** – this is called **chemoradiotherapy**. **Chemotherapy** may be given to try to cure the cancer or to prolong life and control symptoms (**palliative care**) (*Vansteenkiste et al., 2013; Vansteenkiste et al., 2014; Novello et al., 2016*).
- **Targeted therapies** and **antiangiogenic therapies** are drugs that block specific signalling pathways in cancer cells that encourage them to grow (*Novello et al., 2016*).

- **Radiotherapy** is a type of treatment that uses **ionising radiation**, which damages the DNA of cancerous cells, causing the cells to die. It may be used instead of surgery to try to cure early-stage NSCLC. **Radiotherapy** can be given after **chemotherapy** or concurrently (**chemoradiotherapy**). **Radiotherapy** is also used to control symptoms when the cancer is more advanced or has spread to other parts of the body. There are various different techniques for delivering **radiotherapy**, including **SABR** (when available), a type of external beam radiation therapy that delivers a high dose of radiation specifically to the **tumour** (*Vansteenkiste et al., 2013; Novello et al., 2016*).

Your doctor and specialist nurse can discuss all of the possible treatment options available to you to help you to make an informed decision about the best way forward for you.

The response to any treatment you receive will be assessed regularly to see how effective the treatment is and to check whether the benefits outweigh any side effects that you might experience. Evaluation of response is recommended after 2-3 months of **systemic anti-cancer treatment** for stage IV NSCLC. This relies on repetition of the initial imaging tests that showed the cancer (*Novello et al., 2016*).

Treatment options for early (stage I-II) NSCLC

Early-stage NSCLC that is confined to the lung may be curable with surgery

Surgery is the main treatment approach for **early-stage NSCLC** (Vansteenkiste et al., 2013; Vansteenkiste et al., 2014). This involves removing the cancer and some of the nearby **lymph nodes** in the chest. The number of **lymph nodes** removed is dependent on the type of surgery performed. Surgical resection of NSCLC is a major operation and you need to be in good general health to be able to cope with it. The type of operation will either be a **lobectomy** (preferred) or a **wedge (segment) resection** and may be carried out via open surgery or **video-assisted thoracic surgery (VATS)**, depending on the preference of your surgeon (Vansteenkiste et al., 2013; Vansteenkiste et al., 2014).

The **lymph nodes** removed during surgery will be examined under a microscope to check for cancer cells. Knowing if the cancer has spread to the **lymph nodes** also helps your doctors decide if you need further treatment with **adjuvant chemotherapy** or **radiotherapy** (Vansteenkiste et al., 2013; Vansteenkiste et al., 2014).

Adjuvant chemotherapy may be given in patients with stage II NSCLC. Your general health and your postoperative recovery will be taken into account when deciding whether you should be offered **adjuvant chemotherapy**. A combination of two different drugs is preferred (one of them being **cisplatin**), and it is likely that you will be offered 3 or 4 cycles of treatment (Vansteenkiste et al., 2013; Vansteenkiste et al., 2014).

Adjuvant radiotherapy is another option when it has not been possible to completely remove the **tumour** during surgery (Vansteenkiste et al., 2013; Vansteenkiste et al., 2014).

In patients with stage I NSCLC who are unwilling or unable to undergo surgery, **SABR** may be offered. This treatment will be given to you as an outpatient over 3-8 visits. If your **tumour** is larger than 5 cm (stage II NSCLC) and/or is located at the centre of the lung, radical **radiotherapy** using more conventional daily or **accelerated schedules** is preferred (Vansteenkiste et al., 2013; Vansteenkiste et al., 2014).

Treatment of early (stage I-II) NSCLC – summary (Vansteenkiste et al., 2013; Vansteenkiste et al., 2014)

TREATMENT TYPE	PATIENTS	TREATMENT DETAILS	CONSIDERATIONS
Surgery	Stage I or II NSCLC	<ul style="list-style-type: none"> Operation is either: <ul style="list-style-type: none"> Lobectomy: The removal of one of the lobes of the lung (preferred option), or Wedge or segment resection: Only a small amount of the lung is removed (sometimes used for very early NSCLC) Carried out either by open surgery or by VATS 	<ul style="list-style-type: none"> Risks associated with major surgery Recovery time (shorter with VATS) Usually able to go home 3-7 days after surgery Requires post-operative pain control
Adjuvant chemotherapy	Stage II NSCLC, following surgery Stage IB NSCLC following surgery, if primary tumour is >4 cm in size (Not recommended in stage IA NSCLC)	<ul style="list-style-type: none"> A combination of two different drugs usually given intravenously (one of which is cisplatin) Typically, 3-4 cycles of treatment 	<ul style="list-style-type: none"> Need to recover from surgery before starting chemotherapy Pre-existing medical conditions may affect whether you will be suitable for chemotherapy
SABR	Preferred for stage I, if surgery not carried out	<ul style="list-style-type: none"> More precise than conventional radiotherapy; very small areas can be targeted with a high dose Shorter treatment time vs conventional radiotherapy (2-week course) 	<ul style="list-style-type: none"> SABR is associated with low toxicity in patients with COPD and in elderly patients Surgery may be offered afterwards if SABR is not successful or if there are complications
Radical radiotherapy	Tumours >5 cm and/or centrally located Following incomplete surgery	<ul style="list-style-type: none"> Conventional (4-7-week course of treatment of short, daily sessions Monday to Friday) or accelerated schedule (an increased number of treatments delivered over a shorter timeframe) 	

COPD, chronic obstructive pulmonary disease; NSCLC, non-small-cell lung cancer; SABR, stereotactic ablative radiotherapy; VATS, video-assisted thoracic surgery

Treatment options for locally advanced (stage III) NSCLC

Treatment for locally advanced disease is likely to involve different types of therapy

Locally advanced NSCLC represents a very diverse disease (see stages IIIA and IIIB in the AJCC/UICC staging system table) and so it is not possible to recommend a 'one size fits all' approach to treatment. Some patients with stage III NSCLC have a **tumour** that is considered **resectable**, i.e. your doctor/surgeon thinks that it can be completely removed by surgery either straight away or after a course of **chemotherapy** (with or without **radiotherapy**). On the other hand, some patients with stage III NSCLC have a **tumour** that is considered **unresectable**, i.e. surgery is not possible due to the size/location of the **tumour** and involvement of **lymph nodes** in the middle of the chest. The best approach to treatment of stage III NSCLC is therefore likely to be a combination of various treatment types (surgery, **chemotherapy** and/or **radiotherapy**), called **multimodal therapy** (Vansteenkiste et al., 2013; Eberhardt et al., 2015).

In patients staged with potentially **resectable** stage III NSCLC, treatment options are generally either **induction therapy** with **chemotherapy** or **chemoradiotherapy**, followed by surgery (preferred for those whose **tumour** is likely to be completely removed by **lobectomy**) or **chemoradiotherapy**. In patients with **unresectable** stage III NSCLC, the preferred treatment is **chemoradiotherapy**. Alternatively, sequential **chemotherapy** and then **radiotherapy** may be given to patients who are unable to tolerate **concurrent** treatment (Vansteenkiste et al., 2013; Eberhardt et al., 2015).

Chemotherapy is an integral part of the treatment of stage III NSCLC. Generally, a **cisplatin**-based combination regimen (two different drugs) is offered. You will usually be offered 2–4 cycles, whether **chemotherapy** is given alone or as part of course of **chemoradiotherapy**. In some patients who undergo surgery upfront for NSCLC that is thought to be stage I or II, but found to be stage III during surgery, then adjuvant **chemotherapy** will likely be administered after the surgery (Vansteenkiste et al., 2013; Eberhardt et al., 2015).

When **radiotherapy** is given concurrently with **chemotherapy** for stage III NSCLC, it is given as conventional daily doses and treatment should not exceed 7 weeks. It may be given as an **accelerated schedule** as part of a pre-operative **chemoradiotherapy** course, but any potential advantages to the likely outcome of surgery will need to be weighed up against potential greater toxicity. When given **sequentially**, an **accelerated schedule** of **radiotherapy** may be given, i.e. higher doses over a shorter timeframe (Vansteenkiste et al., 2013; Eberhardt et al., 2015).

Treatment of locally advanced (stage III) NSCLC – summary (Vansteenkiste et al., 2013; Eberhardt et al., 2015)

TREATMENT TYPE	PATIENTS	TREATMENT DETAILS	CONSIDERATIONS
Surgery	Resectable stage III NSCLC	<ul style="list-style-type: none"> Preferred when a complete resection by lobectomy is expected, to spare as much lung tissue as possible May require a pneumonectomy (removal of one lung) in some patients May be offered after an initial course of chemotherapy (+/- radiotherapy) – called induction therapy 	<ul style="list-style-type: none"> Outcome depends on the extent of involvement of the lymph nodes at the centre of the chest – may not be known until after surgery Lung function tests are important before deciding on surgery
	Chemotherapy	<ul style="list-style-type: none"> Intravenous cisplatin-based regimen is preferred (cisplatin-etoposide or cisplatin-vinorelbine) Typically, 2-4 cycles of treatment are given 	<ul style="list-style-type: none"> A carboplatin-based combination may be chosen if you have other medical conditions that could affect how you tolerate chemotherapy
	Resectable stage III NSCLC	<ul style="list-style-type: none"> If your tumour is considered resectable, chemotherapy may be given before surgery as induction therapy (chemotherapy +/- radiotherapy) If you have surgery upfront and it is found that the cancer had spread to lymph nodes in the chest, you may be offered adjuvant chemotherapy 	<ul style="list-style-type: none"> It is likely that you will experience more side effects if chemotherapy is given concurrently with radiotherapy
Radiotherapy	Unresectable stage III NSCLC	<ul style="list-style-type: none"> Delivered concurrently with radiotherapy (preferred) or sequentially (before radiotherapy) if concurrent treatment cannot be tolerated 	
	Resectable stage III NSCLC	<ul style="list-style-type: none"> May be given post-operatively in patients who have had incomplete resection When given pre-operatively concurrently with chemotherapy, may be conventional doses or as an accelerated schedule 	
	Unresectable stage III NSCLC	<ul style="list-style-type: none"> May be given as conventional daily doses as part of a chemoradiotherapy schedule (up to 7 weeks), or sequentially (after chemotherapy) as an accelerated schedule 	

NSCLC, non-small-cell lung cancer

Treatment options for metastatic (stage IV) NSCLC

Chemotherapy is the main treatment for metastatic NSCLC

Metastatic NSCLC is usually considered inoperable. Complete removal of the **tumour(s)** is very unlikely and therefore a chance of cure cannot be offered. However, surgical interventions can relieve symptoms caused by the disease spreading to other parts of the body. Similarly, **radiotherapy** may help control symptoms that arise due to the disease spreading to certain organs, including the brain and bones (*Novello et al., 2016*).

Systemic anticancer treatment is the main treatment for stage IV NSCLC, the aims of which are to improve quality of life and to prolong survival. There are many different types of drugs available and the choice of which drugs are offered will largely depend on your general health and the type of **tumour** that you have (*Novello et al., 2016*).

Intravenous chemotherapy with a two-drug combination (**doublet chemotherapy**) is the main treatment for patients with **metastatic** NSCLC whose cancer does not contain specific modifications to the **EGFR** or **ALK** genes or high levels of the **PD-L1** protein (determined by molecular testing using a **tumour biopsy**). **Doublet chemotherapy** is likely to include a **platinum-based** compound plus either **gemcitabine**, **vinorelbine** or a **taxane**. Addition of **pemetrexed** or the **targeted agent bevacizumab** may also be considered for non-squamous NSCLC. In patients whose general health is poor, single-agent **chemotherapy** with **gemcitabine**, **vinorelbine** or **docetaxel** is another treatment option (*Novello et al., 2016*).

Patients whose **tumours** have **EGFR** mutations or **ALK rearrangements** are best treated with oral **targeted therapies**. **Gefitinib**, **erlotinib** or **afatinib** are options for **EGFR**-mutated **tumours** and **crizotinib** is offered to patients who have an **ALK rearrangement** (*Novello et al., 2016*).

Patients whose tumours express relatively high levels of **PD-L1** protein (determined by molecular testing using a **tumour biopsy**) may receive **first-line immunotherapy** with **pembrolizumab**.

After 4–6 cycles of **doublet chemotherapy**, **maintenance treatment** with **pemetrexed**, with or without **bevacizumab**, may be given to patients in good general health with non-squamous **tumours** to prolong the effect of **first-line chemotherapy** on **tumour** control. **Erlotinib** may be offered as **maintenance treatment** in patients whose **tumours** have **EGFR mutations** (*Novello et al., 2016*).

Further lines of treatment may be offered, depending on the **first-line** treatment received and on the general health of the patient. Treatment options include: **Chemotherapy** (**pemetrexed** or **docetaxel**), **immunotherapy** (**nivolumab** or **pembrolizumab**), **antiangiogenic agents** (**nintedanib** or **ramucirumab**) plus **docetaxel**, and **targeted therapies** (**afatinib** or **erlotinib**). Patients whose **tumours** have **EGFR mutations** who have received **first-line** treatment with **erlotinib**, **gefitinib** or **afatinib**, and who have a confirmed abnormality called a **T790M mutation**, may be treated with **second-line osimertinib**. Patients whose **tumours** have **ALK rearrangements** who have received **first-line** treatment with **crizotinib** may be treated with **second-line ceritinib** or **alectinib** (Novello et al., 2016).

Treatment of metastatic (stage IV) NSCLC – summary (Novello et al., 2016)

TREATMENT TYPE	PATIENTS	TREATMENT DETAILS	CONSIDERATIONS
Chemotherapy	EGFR- and ALK-negative tumours <ul style="list-style-type: none"> Good general condition, no other major medical conditions 	First line: <ul style="list-style-type: none"> Intravenous platinum-based regimen preferred (2 drug combination including cisplatin or carboplatin + gemcitabine, vinorelbine or a taxane) Pemetrexed may be incorporated into the treatment regimen in non-squamous histology 4-6 cycles (may be offered maintenance treatment with single agent pemetrexed after 4 cycles) Second line: <ul style="list-style-type: none"> Pemetrexed (non-squamous type) or docetaxel 	<ul style="list-style-type: none"> Response to platinum-based therapy, toxicity and patient's general health after initial treatment needs to be considered when deciding upon maintenance treatment Patients with a very poor general condition are not suitable for chemotherapy; best supportive care is the only treatment
	<ul style="list-style-type: none"> Less fit patients/elderly 	First line: <ul style="list-style-type: none"> Carboplatin-based regimen preferred; may be offered single-agent treatment with gemcitabine, vinorelbine or docetaxel 	
Targeted therapy	EGFR mutation	First line: <ul style="list-style-type: none"> Gefitinib, erlotinib or afatinib Second line: <ul style="list-style-type: none"> Osimertinib 	<ul style="list-style-type: none"> As most targeted agents are generally well tolerated, they may also be offered to patients with a moderate/poor general condition
	ALK rearrangement	First line: <ul style="list-style-type: none"> Crizotinib Second line: <ul style="list-style-type: none"> Ceritinib Alectinib 	
	Targeted therapy in tumours without specific mutations	First line: <ul style="list-style-type: none"> Intravenous bevacizumab may be added to a platinum-based regimen (non-squamous type) in patients in good general condition Necitumumab + gemcitabine/cisplatin is an option for SCC expressing EGFR Second line: <ul style="list-style-type: none"> Erlotinib, nintedanib + docetaxel (adencocarcinoma), ramucirumab + docetaxel, afatinib 	

continued overleaf

TREATMENT TYPE	PATIENTS	TREATMENT DETAILS	CONSIDERATIONS
Immunotherapy	EGFR- and ALK-negative tumours <ul style="list-style-type: none"> Good general condition, no other major medical conditions 	First line: <ul style="list-style-type: none"> Pembrolizumab (in patients with tumours strongly positive for PD-L1) Second line: <ul style="list-style-type: none"> Nivolumab or pembrolizumab (in patients with tumours positive for PD-L1) 	
Surgery	Can be used for relief of symptoms caused by cancer spreading	Minimally-invasive procedures can be helpful, e.g. placement of a stent to alleviate obstruction of the airways	
Radiotherapy	Can be used for relief of symptoms caused by cancer spreading	<ul style="list-style-type: none"> Radiotherapy can achieve symptom control for bone and brain metastases It can also relieve symptoms caused by airway obstruction 	

ALK, anaplastic lymphoma kinase; EGFR, epidermal growth factor; NSCLC, non-small-cell lung cancer; PD-L1, programmed death-ligand 1; SCC, squamous cell carcinoma

Oligometastatic disease

When the cancer has spread beyond the site at which it started but is not yet widely **metastatic**, it is called **oligometastatic disease**. If you have **synchronous oligometastases** diagnosed within 1 month of your **primary tumour**, it may be possible to achieve long-term disease-free survival following **chemotherapy** and radical local treatment, such as high-dose **radiotherapy** or surgery; inclusion in a suitable clinical trial may be advised by your doctor (*Novello et al., 2016*). Similarly, if you have a limited number of **metachronous oligometastases** that appear following treatment for your **primary tumour**, you may be offered treatment with high-dose **radiotherapy** or surgery (*Novello et al., 2016*).

Clinical trials

Your doctor may ask you whether you would like to take part in a **clinical trial**. This is a research study conducted with patients in order to (*ClinicalTrials.gov, 2017*):

- Test new treatments
- Look at new combinations of existing treatments, or change the way they are given to make them more effective or reduce side effects
- Compare the effectiveness of drugs used to control symptoms
- Find out how cancer treatments work.

Clinical trials help to improve knowledge about cancer and develop new treatments, and there can be many benefits to taking part. You would be carefully monitored during and after the study, and the new treatment may offer benefits over existing therapies. It's important to bear in mind, however, that some new treatments are found not to be as good as existing treatments or to have side effects that outweigh the benefits (*ClinicalTrials.gov, 2017*).

Clinical trials help to improve knowledge about diseases and develop new treatments – there can be many benefits to taking part

You have the right to accept or refuse participation in a **clinical trial** without any consequences for the quality of your treatment. If your doctor does not ask you about taking part in a **clinical trial** and you want to find out more about this option, you can ask your doctor if there is a trial for your type of cancer taking place nearby (*ClinicalTrials.gov, 2017*).

Supportive care

There is a range of other therapies available that can also help with the management of NSCLC. These include bone modifying agents (e.g. zoledronic acid and denosumab, used to reduce the occurrence of fractures commonly associated with the presence of bone **metastases**), **stents** (for relieving major airway obstructions that can cause **dyspnoea**), pain management and nutritional support (*Novello et al., 2016*). Generally, early **supportive care** is recommended in parallel with treatments for the cancer itself: it may improve your quality of life and mood and lessen the need for aggressive treatment (*Novello et al., 2016*).

What are the possible side effects of treatment?

As with any medical treatment, you may experience side effects from your anti-cancer treatment. The most common side effects for each type of treatment are highlighted below, along with some information on how they can be managed. You may experience side effects other than those discussed here. It is important to talk to your doctor or **nurse specialist** about any potential side effects that you are concerned about.

Fatigue is very common in patients undergoing cancer treatment, and can result from either the cancer itself or the treatments. Your doctor or nurse can provide you with strategies to limit the impact of **fatigue**, including getting enough sleep, eating healthily and staying active (*Cancer.Net, 2016c*)

It is important to talk to your doctor about any treatment-related side effects that you are concerned about

Surgery

Side effects following cancer surgery vary depending on the location and type of the surgery and your general health (*Cancer.Net, 2016a*). Common side effects following lung **resection** are summarised in the table.

POSSIBLE SIDE EFFECT	HOW THE SIDE EFFECT MAY BE MANAGED
Pain	Pain or discomfort following surgery is common and can usually be controlled using pain-relief medication. Always let your doctor or nurse know if you are in pain, so they can treat it as soon as possible (<i>Macmillan, 2015a</i>)
Infection	You will be taught how to lower the risk of infection occurring. Signs of infection include redness, warmth, increased pain and weeping from around the wound. If you notice any of these signs, contact your nurse or doctor (<i>Cancer.Net, 2016a</i>)
Prolonged air leak	Air leak is a natural occurrence after lung resection but its prolongation to over 7 days increases the risks of other complications. Your surgeon will take precautions to minimise the risk of prolonged air leak (<i>Ziarnik et al., 2015</i>)
Pneumonia	The risk of pneumonia can be decreased by following advice provided by your doctor, e.g. you should perform any recommended physiotherapy exercises (e.g. coughing), start walking/moving about as soon as possible after surgery and refrain from smoking. If pneumonia occurs, then it can usually be treated with an antibiotic (<i>Ziarnik et al., 2015</i>)

Common side effects of lung cancer surgery and how they can be managed

Radiotherapy

For some patients, **radiotherapy** causes few or no side effects; for others, the side effects can be severe. Side effects occur because radiation therapy can damage healthy tissues near the treatment area. The side effects will depend upon the location of the treatment area, the radiation dose and your general health. Usually, side effects start to appear after 2 or 3 weeks of treatment, and resolve a few weeks following the final treatment (*Cancer.Net, 2016b*).

POSSIBLE SIDE EFFECT	HOW THE SIDE EFFECT MAY BE MANAGED
Skin damage (e.g. dryness, itching, blistering, or peeling)	These side effects usually go away a few weeks after treatment has finished. If skin damage becomes a serious problem, then your doctor may change your treatment plan (<i>Cancer.Net, 2016b</i>)
Oesophagitis	After 2–3 weeks of radiotherapy to the chest, you may have difficulty swallowing, heartburn or indigestion. This is because radiotherapy can cause inflammation in the oesophagus . Your doctor or nurse will be able to advise you on how to cope with these symptoms and may prescribe medicines to help (<i>Macmillan, 2015b</i>)
Radiation pneumonitis (cough, fever and fullness of chest)	Patients receiving radiotherapy to the chest may develop a condition called radiation pneumonitis . This generally appears between 2 weeks and 6 months following radiotherapy but is usually temporary. Tell your doctor or nurse if you experience any of the signs of radiation pneumonitis (<i>Cancer.Net, 2016b</i>)

*Common side effects of **radiotherapy** used to treat lung cancer and how they can be managed*

Chemotherapy

Side effects from **chemotherapy** vary depending upon the drugs and the doses used – you may get some of those listed below but you are very unlikely to get all of them. Patients who receive a combination of different **chemotherapy** drugs are likely to experience more side effects than those who receive a single **chemotherapy** drug. The main areas of the body affected by **chemotherapy** are those where new cells are being quickly made and replaced (**bone marrow, hair follicles**, the digestive system, the lining of your mouth). Reductions in your levels of **neutrophils** (a type of white blood cell) can lead to **neutropenia**, which will make you more susceptible to infections. Some **chemotherapy** drugs can affect fertility – if you are worried about this, speak to your doctor before treatment starts. Most side effects of **chemotherapy** are temporary and can be controlled with drugs or lifestyle changes – your doctor or nurse will help you to manage them (Macmillan, 2016a).

CHEMOTHERAPY DRUG	POSSIBLE SIDE EFFECT	HOW THE SIDE EFFECTS MAY BE MANAGED
Cisplatin (Macmillan, 2016b)	<ul style="list-style-type: none">• Increased risk of infection• Neutropenia• Thrombocytopenia• Anaemia• Nausea/vomiting• Anorexia• Changes in kidney function• Tinnitus/changes in hearing• Peripheral neuropathy• Fatigue• Taste changes• Diarrhoea• Decreased fertility• Increased risk of thrombosis	<ul style="list-style-type: none">• Your blood cell counts will be monitored frequently throughout your treatment in order to detect any neutropenia, anaemia or thrombocytopenia – your doctor may adjust your treatment according to test results, and will advise you on how to prevent infections.• Effects on the gastrointestinal system (nausea, vomiting, diarrhoea, taste changes) may result in loss of appetite (anorexia). Your doctor will be able to help you to prevent or manage these side effects.• Report any signs of peripheral neuropathy (tingling or numbness in your hands or feet) to your doctor, who will help you to manage this side effect.• You will have tests before and during treatment to check how well your kidneys are functioning. You will be asked to drink plenty of fluids to prevent your kidneys from becoming damaged.• Tell your doctor if you notice any changes in your hearing or experience tinnitus. Changes in hearing are usually temporary but can occasionally be permanent.
Carboplatin (Macmillan, 2015c)	<ul style="list-style-type: none">• Neutropenia• Thrombocytopenia• Anaemia• Increased risk of infection• Nausea• Vomiting• Constipation• Fatigue• Renal (kidney) toxicity• Hepatic (liver) toxicity	<ul style="list-style-type: none">• Your blood cell counts will be monitored frequently throughout your treatment in order to detect any neutropenia, anaemia or thrombocytopenia – your doctor may adjust your treatment according to test results, and will advise you on how to prevent infections.• Your doctor will be able to help you prevent or manage any nausea, vomiting or constipation.• You will have tests before and during treatment to check how well your kidneys and liver are functioning, and you will be asked to drink plenty of fluids to prevent your kidneys from becoming damaged.

continued overleaf

CHEMOTHERAPY DRUG	POSSIBLE SIDE EFFECT	HOW THE SIDE EFFECTS MAY BE MANAGED
Docetaxel (Taxotere SPC, 2005)	<ul style="list-style-type: none"> • Neutropenia • Increased risk of infections • Anaemia • Thrombocytopenia • Peripheral neuropathy • Nausea • Vomiting • Diarrhoea • Stomatitis • Anorexia • Asthenia • Skin reaction • Oedema • Alopecia 	<ul style="list-style-type: none"> • Your blood cell counts will be monitored frequently throughout your treatment in order to detect any neutropenia, anaemia or thrombocytopenia – your doctor may adjust your treatment according to test results, and will advise you on how to prevent infections. • Report any signs of peripheral neuropathy (tingling or numbness in your hands or feet) to your doctor, who will help you to manage this side effect. • Effects on the gastrointestinal system (nausea, vomiting, diarrhoea) and stomatitis may result in loss of appetite (anorexia) or feelings of weakness (asthenia). Your doctor will be able to help you to prevent or manage these side effects. • Let your doctor know if you experience any skin reactions or fluid retention/swelling (oedema) – they will help you to manage these side effects. • Alopecia can be upsetting for many patients; your doctor will provide you with information on how to cope with this side effect.
Etoposide (Vepesid SPC, 2016)	<ul style="list-style-type: none"> • Neutropenia • Anaemia • Leukopenia • Thrombocytopenia • Constipation • Nausea • Vomiting • Anorexia • Asthenia • Changes in liver function • Alopecia 	<ul style="list-style-type: none"> • Your blood cell counts will be monitored frequently throughout your treatment in order to detect any neutropenia, anaemia, thrombocytopenia or leukopenia – your doctor may adjust your treatment according to test results, and will advise you on how to prevent infections. • Effects on the gastrointestinal system (constipation, nausea, vomiting) may result in loss of appetite (anorexia) or feelings of fatigue/asthenia. Your doctor will be able to help you to prevent or manage these side effects. • You will have tests before and during treatment to check how well your liver is functioning. • Alopecia can be upsetting for many patients; your doctor will provide you with information on how to cope with this side effect.

CHEMOTHERAPY DRUG	POSSIBLE SIDE EFFECT	HOW THE SIDE EFFECTS MAY BE MANAGED
Paclitaxel (Abraxane SPC, 2013)	<ul style="list-style-type: none"> • Neutropenia • Anaemia • Leukopenia • Thrombocytopenia • Lymphopenia • Fever • Nausea • Diarrhoea • Vomiting • Constipation • Stomatitis • Anorexia • Fatigue • Asthenia • Rash • Arthralgia • Myalgia • Peripheral neuropathy • Alopecia 	<ul style="list-style-type: none"> • Your blood cell counts will be monitored frequently throughout your treatment in order to detect any neutropenia, anaemia, leukopenia, thrombocytopenia or lymphopenia – your doctor may adjust your treatment according to test results, and will advise you on how to prevent infections. Report any fever to your doctor, as this may be a sign of infection. • Effects on the gastrointestinal system (nausea, vomiting, diarrhoea, constipation, stomatitis) may result in loss of appetite (anorexia) or feelings of fatigue/asthenia. Your doctor will be able to help you to prevent or manage these side effects. • Let your doctor know if you experience arthralgia, myalgia or rash and they will help you to manage these side effects. • Report any signs of peripheral neuropathy (tingling or numbness in your hands or feet) to your doctor, who will help you to manage this side effect. • Alopecia can be upsetting for many patients; your doctor will provide you with information on how to cope with this side effect.
Pemetrexed (Alimta SPC, 2017)	<ul style="list-style-type: none"> • Neutropenia • Anaemia • Leukopenia • Stomatitis • Pharyngitis • Nausea • Anorexia • Fatigue • Rash 	<ul style="list-style-type: none"> • Your blood cell counts will be monitored frequently throughout your treatment in order to detect any neutropenia, anaemia, or leukopenia – your doctor may adjust your treatment according to test results, and will advise you on how to prevent infections. • Effects on the gastrointestinal system (stomatitis, pharyngitis, nausea) may result in loss of appetite (anorexia). Your doctor will be able to help you to prevent or manage these side effects. • Let your doctor know if you develop a rash – they will help you to manage this side effect.

CHEMOTHERAPY DRUG	POSSIBLE SIDE EFFECT	HOW THE SIDE EFFECTS MAY BE MANAGED
Vinorelbine (Vinorelbine SPC, 2014)	<ul style="list-style-type: none"> • Neutropenia • Anaemia • Neurological disorders • Stomatitis • Nausea • Vomiting • Constipation • Oesophagitis • Skin reactions • Alopecia 	<ul style="list-style-type: none"> • Your blood cell counts will be monitored frequently throughout your treatment in order to detect any neutropenia or anaemia – your doctor may adjust your treatment according to test results, and will advise you on how to prevent infections. • Report any signs of neurological disorders (e.g. loss of reflexes, weakness of the legs and feet) to your doctor, who will decide how to manage these side effects. • Your doctor will be able to help you to prevent or manage any effects on the gastrointestinal system (stomatitis, nausea, vomiting, constipation, oesophagitis). • Let your doctor know if you experience any burning or skin changes at the injection site, so that they can decide how to manage these. • Alopecia can be upsetting for many patients; your doctor will provide you with information on how to cope with this side effect.

Very common side effects with chemotherapy (used as single drugs) in the treatment of NSCLC. The most recent Summary of Product Characteristics (SPCs) for individual drugs can be located at: <http://www.ema.europa.eu/ema/>.

Targeted therapies, antiangiogenic therapies and immunotherapies

Common side effects in patients treated with **targeted therapies**, **antiangiogenic therapies** or **immunotherapy** include effects on the **gastrointestinal system** (e.g. diarrhoea, vomiting, nausea), skin problems (e.g. rash, dry skin, nail changes, discolouration) and **hypertension** (high blood pressure). Many of the side effects from **targeted therapies** can be effectively prevented or managed effectively. Always tell your doctor or nurse as soon as possible if you notice any side effects from taking a **targeted therapy**, **antiangiogenic therapy** or **immunotherapy**.

THERAPY	POSSIBLE SIDE EFFECT	HOW THE SIDE EFFECTS MAY BE MANAGED
Alectinib (Alecensa SPC, 2017)	<ul style="list-style-type: none">• Nausea• Constipation• Oedema• Myalgia	<ul style="list-style-type: none">• Report any nausea or constipation to your doctor, who will be able to help you to prevent or manage these side effects.• Let your doctor know if you develop any oedema (fluid retention) or myalgia (muscle pain) – they will help you to manage these side effects.
Bevacizumab (Avastin SPC, 2016)	<ul style="list-style-type: none">• Neutropenia• Leukopenia• Thrombocytopenia• Peripheral neuropathy• Wound healing complications• Bleeding disorders• Hypertension• Stomatitis• Constipation• Diarrhoea• Nausea• Vomiting• Anorexia• Skin reactions• Dysguesia• Fatigue• Dysarthria• Headache• Watery eyes• Dyspnoea• Rhinitis• Arthralgia	<ul style="list-style-type: none">• Your blood cell counts will be monitored frequently throughout your treatment in order to detect any neutropenia, leukopenia or thrombocytopenia – your doctor may adjust your treatment according to test results, and will advise you on how to prevent infections.• Report any signs of peripheral neuropathy (tingling or numbness in your hands or feet) to your doctor, who will help you to manage this side effect.• Any treatment will be delayed until wounds have healed satisfactorily.• Your blood pressure will be monitored throughout treatment and any hypertension will be managed appropriately.• Effects on the gastrointestinal system (stomatitis, constipation, diarrhoea, nausea, vomiting) and dysguesia (taste changes) may result in loss of appetite (anorexia). Your doctor will be able to help you to prevent or manage these side effects.• Let your doctor know if you develop any skin reactions (e.g. rash, dry skin, discolouration) – they will help you to manage these side effects.• Report any other side effects, including changes in vision, dyspnoea (breathlessness), dysarthria (difficulty with speech), arthralgia (painful joints) or headache to your doctor, who will help you to manage these side effects.

THERAPY	POSSIBLE SIDE EFFECT	HOW THE SIDE EFFECTS MAY BE MANAGED
Gefitinib (Iressa SPC, 2016)	<ul style="list-style-type: none"> • Diarrhoea • Anorexia • Asthenia • Changes in liver function • Skin reactions 	<ul style="list-style-type: none"> • Diarrhoea may result in loss of appetite (anorexia) and asthenia (weakness). Your doctor will be able to help you to prevent or manage these side effects. • You will have tests before and during treatment to check how well your liver is functioning. • Let your doctor know if you develop any skin reactions (e.g. rash, acne, dry skin, itchiness) – they will help you to manage these side effects.
Erlotinib (Tarceva SPC, 2017)	<ul style="list-style-type: none"> • Increased risk of infection • Diarrhoea • Nausea • Vomiting • Stomatitis • Anorexia • Fatigue • Dry eyes • Conjunctivitis • Dyspnoea • Cough • Rash 	<ul style="list-style-type: none"> • Your doctor will advise you on how to prevent infections. • Effects on the gastrointestinal system (diarrhoea, nausea, vomiting, stomatitis) may result in loss of appetite (anorexia). Your doctor will be able to help you to prevent or manage these side effects. • Let your doctor know if you develop any problems with your eyes (e.g. dry eyes, conjunctivitis), experience increased dyspnoea (breathlessness) or cough, or develop a rash – they will help you to manage these side effects.
Afatinib (Giotrif SPC, 2016)	<ul style="list-style-type: none"> • Diarrhoea • Nausea • Vomiting • Stomatitis • Decreased appetite • Epistaxis • Skin reactions (rash, acne, dry skin, itchiness) • Nail disorders 	<ul style="list-style-type: none"> • Effects on the gastrointestinal system (diarrhoea, nausea, vomiting, stomatitis) may result in loss of appetite (anorexia). Your doctor will be able to help you to prevent or manage these side effects. • Let your doctor know if you experience epistaxis (nose bleeds) – they will help you to manage this side effect. • Report any skin reactions or nail changes to your doctor – they will help you to manage these side effects.
Crizotinib (Xalkori SPC, 2017)	<ul style="list-style-type: none"> • Neutropenia • Anaemia • Leukopenia • Peripheral neuropathy • Changes in liver function • Vomiting • Diarrhoea • Nausea • Constipation • Dysgeusia • Fatigue • Impaired vision • Bradycardia • Dizziness • Oedema • Rash 	<ul style="list-style-type: none"> • Your blood cell counts will be monitored frequently throughout your treatment in order to detect any neutropenia, anaemia or leukopenia – your doctor may adjust your treatment according to test results, and will advise you on how to prevent infections. • Report any signs of peripheral neuropathy (tingling or numbness in your hands or feet) to your doctor, who will help you to manage this side effect. • You will have tests before and during treatment to check how well your liver is functioning. • If you experience diarrhoea, nausea, vomiting, constipation, or changes in your sense of taste (dysgeusia), your doctor will be able to help you to prevent or manage these side effects. • Let your doctor know if you develop any problems with your eyes, experience dizziness, oedema (fluid retention) or develop a rash – they will help you to manage these side effects.

Non-small-cell lung cancer

THERAPY	POSSIBLE SIDE EFFECT	HOW THE SIDE EFFECTS MAY BE MANAGED
Ramucirumab^a (Cyramza SPC, 2016)	<ul style="list-style-type: none"> • Neutropenia • Thrombocytopenia • Hypertension • Epistaxis • Stomatitis • Fatigue/asthenia • Oedema 	<ul style="list-style-type: none"> • Your blood cell counts will be monitored frequently throughout your treatment in order to detect any neutropenia or thrombocytopenia – your doctor may adjust your treatment according to test results, and will advise you on how to prevent infections. • Your blood pressure will be monitored throughout treatment and any hypertension will be managed appropriately. • Let your doctor know if you experience a sore mouth or lips, or oedema (fluid retention), your doctor will be able to help you to prevent or manage these side effects.
Nintedanib^a (Vartegaf SPC, 2016)	<ul style="list-style-type: none"> • Neutropenia • Peripheral neuropathy • Diarrhoea • Vomiting • Nausea • Mucositis • Stomatitis • Changes in liver function • Rash 	<ul style="list-style-type: none"> • Your blood cell counts will be monitored frequently throughout your treatment in order to detect any neutropenia – your doctor may adjust your treatment according to test results, and will advise you on how to prevent infections. • Report any signs of peripheral neuropathy (tingling or numbness in your hands or feet) to your doctor, who will help you to manage this side effect. • If you experience diarrhoea, nausea, vomiting, a sore mouth or lips, your doctor will be able to help you to prevent or manage these side effects. • You will have tests before and during treatment to check how well your liver is functioning. • Let your doctor know if you develop any rash – they will help you to manage this side effect.
Necitumumab^b (Portrazza SPC, 2016)	<ul style="list-style-type: none"> • Vomiting • Stomatitis • Weight loss • Altered levels of minerals and salts (hypocalcaemia, hypophosphataemia, hypokalaemia, hypomagnesaemia) • Skin reactions • Fever 	<ul style="list-style-type: none"> • If you experience vomiting or a sore mouth or lips, your doctor will be able to help you to prevent or manage these side effects, and limit any weight loss. • Your body's levels of minerals and salts will be monitored during your treatment – your treatment may be adapted according to any changes. • Let your doctor know if you experience any skin reactions (e.g. rash, dry skin, discolouration), your doctor will be able to help you to prevent or manage these side effects.
Osimertinib (Tagrisso SPC, 2017)	<ul style="list-style-type: none"> • Neutropenia • Leukopenia • Thrombocytopenia • Diarrhoea • Stomatitis • Skin reactions (rash, dry skin, itchiness) • Nail disorders 	<ul style="list-style-type: none"> • Your blood cell counts will be monitored frequently throughout your treatment in order to detect any neutropenia, leukopenia or thrombocytopenia – your doctor may adjust your treatment according to test results, and will advise you on how to prevent infections. • If you experience diarrhoea or a sore mouth or lips, your doctor will be able to help you to prevent or manage these side effects. • Report any skin reactions or nail changes to your doctor – they will help you to manage these side effects.

THERAPY	POSSIBLE SIDE EFFECT	HOW THE SIDE EFFECTS MAY BE MANAGED
Ceritinib (Zykadia SPC, 2016)	<ul style="list-style-type: none"> • Anaemia • Changes in liver function • Diarrhoea • Nausea • Vomiting • Constipation • Dyspepsia, acid reflux, dysphagia • Decreased appetite • Fatigue • Rash 	<ul style="list-style-type: none"> • Your blood cell counts will be monitored frequently throughout your treatment in order to detect any anaemia – your doctor may adjust your treatment according to test results. • You will have tests before and during treatment to check how well your liver is functioning. • If you experience diarrhoea, nausea, vomiting, constipation, indigestion, heartburn or problems swallowing, your doctor will be able to help you to prevent or manage these side effects. • Report any rashes to your doctor – they will help you to manage this side effect.
Nivolumab (Opdivo SPC, 2015)	<ul style="list-style-type: none"> • Neutropenia • Lymphopenia • Leukopenia • Thrombocytopenia • Anaemia • Changes in liver function • Diarrhoea • Nausea • Fatigue • Altered levels of minerals and salts (hypercalcaemia, hyperkalaemia, hypokalaemia, hypomagnesaemia, hyponatraemia) • Rash • Pruritus 	<ul style="list-style-type: none"> • Your blood cell counts will be monitored frequently throughout your treatment in order to detect any neutropenia, lymphopenia, leukopenia, anaemia or thrombocytopenia – your doctor may adjust your treatment according to test results, and will advise you on how to prevent infections. • You will have tests before and during treatment to check how well your liver is functioning. • Your doctor will be able to help you to prevent or manage any diarrhoea or nausea. • Your body's levels of minerals and salts will be measured during your treatment – your treatment may be adapted according to any changes. • Let your doctor know if you experience any skin rash or itchiness – they will be able to help you to prevent or manage these side effects.
Pembrolizumab (Keytruda SPC, 2015)	<ul style="list-style-type: none"> • Diarrhoea • Nausea • Rash • Pruritus • Arthralgia • Fatigue 	<ul style="list-style-type: none"> • Your doctor will be able to help you to prevent or manage any diarrhoea or nausea. • Let your doctor know if you experience any skin rash or itchiness or joint pain – they will be able to help you to prevent or manage these side effects.

Very common side effects with targeted therapy and antiangiogenic therapy in the treatment of NSCLC. The most recent Summary of Product Characteristics (SPCs) for individual drugs can be located at: <http://www.ema.europa.eu/ema/>.

^aIn combination with docetaxel chemotherapy; ^bVery common side effects reported when administered in combination with gemcitabine and cisplatin.

What happens after my treatment has finished?

Follow-up appointments

You will be able to discuss any concerns you have at your follow-up appointments

After your treatment has finished, your doctor will arrange follow-up appointments. You will have regular chest **x-rays** and/or **CT** scans to check that there are no further **tumours**. Your doctor will also evaluate any treatment complications or side effects related to surgery, **radiotherapy** and/or **chemotherapy**. The frequency of these appointments will be tailored to your situation, and will depend on the stage of the cancer when you were initially diagnosed and the treatment that you have had (*Vansteenkiste et al., 2013; Novello et al., 2016*).

Recommendations

- After surgery for stage I-III NSCLC, you should be seen every 3-6 months for the first 2-3 years and then yearly after that (*Vansteenkiste et al., 2013*).
- After **SABR**, you may have a **CT** scan every 6 months, particularly if you are suitable for salvage treatment should there be any complications (*Vansteenkiste et al., 2014*).
- After treatment for **metastatic** disease, depending on your suitability for further treatment, your doctor will see you every 6-12 weeks so that **second-line** therapy can be started promptly, if needed (*Novello et al., 2016*).
- If you have had multi-modality treatment for stage III disease you are likely to have brain scans to monitor for the development of brain **metastases**, for which you may be offered treatment (*Eberhardt et al., 2015*).

What if I need more treatment?

Cancer that comes back is called a recurrence. The treatment that you will be offered depends on the extent of the recurrence. When the **tumour** comes back as a recurrence at a single site, you may be offered treatment such as surgical removal or **radiotherapy**. However, this approach is limited to a very small group of patients. Recurrent **tumours** are normally regarded as **metastatic** cancers and you can usually have further **chemotherapy** with different drugs. Sometimes, **targeted therapy** drugs are given with **chemotherapy** (see section 'Treatment options for metastatic (stage IV) NSCLC' for more information).

In some cases, a repeated **biopsy** of the **tumour** may be carried out as it may result in a change to the treatment decision. This may be particularly true if you have been cancer-free for some time after surgical resection. Where available, patients who were previously treated for NSCLC with an **EGFR**-activating **mutation** may undergo a **liquid biopsy** to detect any T790M **mutation** (also called plasma **EGFR** mutational analysis). This will involve providing a small blood sample for analysis. Re-**biopsy** may be useful to differentiate between disease

recurrence and a new **primary lung tumour** (if the recurrence is detected in the lung) to ascertain the type of **tumour** or to repeat the **EGFR mutation** test if a non-squamous cancer is detected (*Vansteenkiste et al., 2013; Vansteenkiste et al., 2014; Eberhardt et al., 2015; Novello et al., 2016*).

Looking after your health

After you have had treatment for NSCLC, you may feel very tired and emotional. It is important to take good care of yourself and get the support that you need.

- **Stop smoking:** If you are a smoker, it is important to stop smoking as soon as you can as it may reduce the risk of disease recurrence (*Vansteenkiste et al., 2013; Vansteenkiste et al., 2014; Eberhardt et al., 2015; Novello et al., 2016*). Your doctor and nurse can offer help with stopping smoking.
- **Take plenty of rest when you need it:** Give your body time to recover and make sure you rest as much as you can. Complementary therapies, such as aromatherapy, may help you relax and cope better with side effects. Your hospital may offer complementary therapy; ask your doctor for details.
- **Eat well and keep active:** Eating a healthy diet and keeping active can help improve your fitness. It is important to start slowly, with gentle walking, and build up as you start to feel better.

Emotional support

It is common to be overwhelmed by your feelings when you have been diagnosed with cancer and when you have been through treatment. If you feel anxious or depressed, talk to your doctor or nurse – they can refer you to a specialist counsellor or psychologist who has experience of dealing with emotional problems of people dealing with cancer. It may also help to join a support group so that you can talk to other people who understand exactly what you are going through.



Support groups

In Europe, there are some lung cancer patient advocacy groups, which help patients and their families to navigate the lung cancer landscape. They can be local, national or international, and they work to ensure patients receive appropriate and timely care and education. These groups can provide you with the tools you may need to help you better understand your disease, and to learn how to cope with it, living the best quality of life that you can.

You can access information from the following organisations:

- **Global Lung Cancer Coalition (GLCC):** www.lungcancercoalition.org
- **Lung Cancer Europe (LuCE):** www.lungcancereurope.eu
- **Women Against Lung Cancer in Europe (WALCE) educational booklets:**
www.womenagainstlungcancer.eu/?lang=en

References

- American Joint Commission on Cancer (AJCC) Cancer Staging Manual. Springer International Publishing. Editor Amin MB, *et al.* 2016.
- Bailey-Wilson JE, Amos CI, Pinney SM, *et al.* A major lung cancer susceptibility locus maps to chromosome 6q23-25. *Am J Hum Genet* 2004;75(3):460-474.
- Cancer.Net. 2016a. Side effects of surgery. Available from: <http://www.cancer.net/navigating-cancer-care/how-cancer-treated/surgery/side-effects-surgery>. Accessed 27th January 2017.
- Cancer.Net. 2016b. Side effects of radiation therapy. Available from: <http://www.cancer.net/navigating-cancer-care/how-cancer-treated/radiation-therapy/side-effects-radiation-therapy>. Accessed 27th January 2017.
- Cancer.Net. 2016c. Fatigue. Available from: <http://www.cancer.net/navigating-cancer-care/side-effects/fatigue>. Accessed 9th February 2017.
- ClinicalTrials.gov. Learn about clinical studies. Available from: <https://clinicaltrials.gov/ct2/about-studies/learn>. Accessed 14th February 2017.
- De Leyn P, Lardinois D, Van Schil PE, *et al.* ESTS guidelines for preoperative lymph node staging for non-small cell lung cancer. *Eur J Cardiothorac Surg* 2007;32(1):1-8.
- Eberhardt WEE, De Ruysscher D, Weder W, *et al.* 2nd ESMO Consensus Conference in Lung Cancer: locally advanced stage III non-small-cell lung cancer. *Ann Oncol* 2015;26:1573-1588.
- Ferlay J, Steliarova-Foucher E, Lortet-Tieulent J, *et al.* Cancer incidence and mortality patterns in Europe: estimates for 40 countries in 2012. *Eur J Cancer* 2013;49(6):1374-1403.
- Gould MK, Donington J, Lynch WR, *et al.* Evaluation of individuals with pulmonary nodules: when is it lung cancer? *Chest* 2013; 143(5 Suppl):e93S-120S.
- Macmillan. 2016a. Possible side effects of Chemotherapy. Available from: <http://www.macmillan.org.uk/information-and-support/lung-cancer/non-small-cell-lung-cancer/treating/chemotherapy/side-effects-of-chemotherapy/possible-side-effects.html>. Accessed 9th February 2017.
- Macmillan. 2016b. Cisplatin. Available from: <https://www.macmillan.org.uk/cancerinformation/cancertreatment/treatmenttypes/Chemotherapy/individualdrugs/cisplatin.aspx>. Accessed 27th January 2017.
- Macmillan. 2015a. What happens after surgery. Available from: <http://www.macmillan.org.uk/information-and-support/anal-cancer/treating/surgery/surgery-explained/what-happens-after-surgery.html>. Accessed 9th February 2017.
- Macmillan. 2015b. Possible side effects of radiotherapy. Available from: <http://www.macmillan.org.uk/information-and-support/lung-cancer/non-small-cell-lung-cancer/treating/radiotherapy/radiotherapy-explained/possible-side-effects.html#236381>. Accessed: 9th February 2017.

Macmillan. 2015c. Carboplatin. Available from: <https://www.macmillan.org.uk/cancerinformation/cancertreatment/treatmenttypes/chemotherapy/individualdrugs/carboplatin.aspx>. Accessed 27th January 2017.

Malvezzi, M, Carioli G, Bertuccio P, *et al*. European cancer mortality predictions for the year 2016 with focus on leukaemias. *Ann Oncol* 2016;27(4):725–731.

Novello S, Barlea F, Calfano R, *et al*. Metastatic non-small-cell lung cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2016;27(Suppl 5):v1–v27.

Torre LA, Bray F, Siegel RL, *et al*. Global cancer statistics, 2012. *CA Cancer J Clin* 2015;65:87–108.

Vansteenkiste J, De Ruyscher D, Eberhardt EE, *et al*. Early and locally advanced non-small-cell lung cancer (NSCLC): ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2013;24(Suppl 6):vi89–vi98.

Vansteenkiste J, Crinò L, Dooks C, *et al*. 2nd ESMO Consensus Conference on Lung Cancer: early-stage non-small-cell lung cancer consensus on diagnosis, treatment and follow-up. *Ann Oncol* 2014;25:1462–1474.

Ziarnik E, Grogan EL. Post-lobectomy early complications. *Thorac Surg Clin* 2015;25(3):355–364.

GLOSSARY

ACCELERATED SCHEDULE

A higher dose of radiation is given at each treatment and there are fewer total treatments than in a **conventional radiotherapy** schedule. The total amount of radiation given is about the same in each schedule

ADENOCARCINOMA

The most common type of lung cancer; it develops from mucus-producing cells that line the airways

ADRENAL GLANDS

Glands in the body that produce hormones, such as adrenaline and steroids. They are located above the kidneys

ADJUVANT (CHEMOTHERAPY OR RADIOTHERAPY)

Additional treatment given after the primary treatment to reduce the chance of the cancer coming back; usually refers to **radiotherapy** and/or **chemotherapy** after surgery

AFATINIB

A type of **targeted therapy** called a **tyrosine kinase inhibitor**, which works by blocking signals within cancer cells and stopping the action of **epidermal growth factor receptor**, causing cancer cells to die. It is administered as a once daily tablet

AIR LEAK

When air escapes from the airways (**bronchioles**, **alveoli**) into the parts of the lung where air is not usually present

ALECTINIB

A type of **targeted therapy** called a **tyrosine kinase inhibitor**, which works by blocking a protein called **anaplastic lymphoma kinase**. It only works in cancer cells with an abnormal version of this protein. It is administered twice-daily as oral capsules

ANAPLASTIC LYMPHOMA KINASE REARRANGEMENTS (ALK)

Anaplastic lymphoma kinase is a cell surface protein. Rearrangement of the ALK gene is an abnormality found in some cancer cells, including NSCLC

ALOPECIA

Hair loss

ALVEOLI

Tiny air sacs within the lungs that allow oxygen and carbon dioxide to move between the lungs and bloodstream

ANAEMIA

A condition characterised by the shortage of red blood cells or haemoglobin (a protein in red blood cells that carries oxygen throughout the body)

ANOREXIA

A lack or loss of appetite

ANTIANGIOGENIC THERAPY

A type of therapy that interferes in the growth and survival of new **blood vessels** (angiogenesis), which plays a critical role in the growth and spread of cancer

ANTIBIOTIC

A type of drug used to treat and prevent bacterial infections

ARSENIC

A naturally occurring substance that has been widely used in some industries (copper or lead smelting; agriculture/pesticides), but has been linked to cancer, including lung cancer

ARTHRALGIA

Pain in a joint(s)

ASBESTOS

A natural, fibrous material that was previously widely used as a building material. Its use is now banned throughout Europe as it is linked to lung diseases, including cancer

ASTHENIA

Abnormal feeling of weakness or lack of energy

BEVACIZUMAB

A type of **targeted therapy** used to treat some cancers, including advanced NSCLC. It is a **monoclonal antibody** that targets **vascular endothelial growth factor** and prevents the cancer cells from developing their own blood supply, thus helping to slow down **tumour** growth

BIOPSY

A medical procedure in which a small sample of cells or tissue is taken for examination under a microscope

BLOOD VESSELS

The structures (tubes) carrying blood through the tissues and organs of the body – they include veins, arteries and capillaries

BONE MARROW

A spongy tissue found inside some bones (e.g. hip and thigh bones). It contains stem cells, which are cells that can develop into the red blood cells, white blood cells or **platelets**

BRADYCARDIA

Abnormally slow heart rate

BRONCHI

The right bronchus and the left bronchus (the **bronchi**) are the two main airways that take air into the lungs

GLOSSARY

BRONCHIOLES

The **bronchi** divide into smaller bronchioles, which then lead to the **alveoli**

BRONCHOSCOPE

A thin, fibre-optic cable that is inserted into the airways (usually through the nose or mouth)

BRONCHOSCOPY

A clinical investigation where your doctor examines your airways using a **bronchoscope**

CARBOPLATIN

A type of **chemotherapy** that is administered through a drip into a vein in your arm or chest

CERITINIB

A type of **targeted therapy** that works by inhibiting a protein called **anaplastic lymphoma kinase**. It is administered as a once-daily capsule to patients who have previously received **crizotinib**

CHEMORADIOTHERAPY

Chemotherapy and **radiotherapy** given together

CHEMOTHERAPY

A type of cancer treatment using medicine that kills the cancer cells by damaging them, so that they cannot reproduce and spread

CHRONIC OBSTRUCTIVE PULMONARY DISEASE (COPD)

A type of lung disease characterised by long-term poor airflow. The main symptoms include shortness of breath and cough

CISPLATIN

A type of **chemotherapy** that is administered through a drip into a vein in your arm or chest

CLINICAL TRIAL

A study that compares the effects of one treatment with another

COMORBIDITIES

Additional diseases or disorders experienced by the patient at the same time

COMPUTED TOMOGRAPHY (CT)

A scan using **x-rays** and a computer to create detailed images of the inside of your body

CONCURRENT

Different types of treatment (e.g. **chemotherapy** and **radiotherapy**) given at the same time

CONJUNCTIVITIS

Inflammation of the membrane that covers the eyeball and lines the eyelid

CONVENTIONAL RADIOTHERAPY

Refers to **radiotherapy** that is given to the **tumour** as a fraction of the complete dose over several sessions – treatment usually consists of a small daily dose over several weeks

CRIZOTINIB

A type of **targeted therapy** called a **tyrosine kinase inhibitor**, which works by blocking a protein called **anaplastic lymphoma kinase**. It only works in cancer cells with an abnormal version of this protein. It is administered as a twice-daily capsule

DIAPHRAGM

The muscle that separates the chest cavity from the abdomen; the diaphragm contracts and relaxes as we breathe in and out

DOCETAXEL

A type of **chemotherapy** that is administered through a drip into a vein in your arm or chest

DOUBLET CHEMOTHERAPY

A combination of two different types of **chemotherapy** administered at the same time

DYSARTHRIA

Difficult or unclear articulation of speech (e.g. slurred, nasal-sounding, hoarse or excessively loud or quiet)

DYSGUESIA

A change in the sense of taste

DYSPEPSIA

The medical term for indigestion

DYSPHAGIA

The medical term for difficulties with swallowing

DYSPNOEA

Shortness of breath

EARLY-STAGE (CANCER)

Cancer that has not spread to the **lymph nodes** or other parts of the body

EPIDERMAL GROWTH FACTOR RECEPTOR (EGFR)

A protein involved in cell growth and division. It is found in abnormally high amounts on the surface of many types of cancer cells

EPISTAXIS

The medical term for a nose bleed

GLOSSARY

ERLOTINIB

A type of **targeted therapy** called a **tyrosine kinase inhibitor**, which works by blocking signals within cancer cells and stopping the action of **epidermal growth factor receptor**, causing cancer cells to die. It is administered as a once-daily tablet

ETOPOSIDE

A type of **chemotherapy** that is administered through a drip into a vein in your arm or chest, or as an oral tablet or capsule

FATIGUE

Overwhelming tiredness

FIRST-LINE

The initial treatment given to a patient

GASTROINTESTINAL SYSTEM

The system of organs responsible for getting food into and out of the body and for making use of food to keep the body healthy – includes the **oesophagus**, stomach and intestines

GEFITINIB

A type of **targeted therapy** called a **tyrosine kinase inhibitor**, which works by blocking signals within cancer cells that tell the cells to grow. It is administered as a once-daily tablet.

GEMCITABINE

A type of **chemotherapy** that is administered through a drip into a vein in your arm or chest

GENERAL ANAESTHETIC

A medication that causes a reversible loss of consciousness

GRADE

Cancer **grade** is based on how different **tumour** cells look from normal cells under a microscope, and on how quickly they grow. The **grade** will be a value between one and three and reflects the aggressiveness of **tumour** cells; the higher the **grade**, the more aggressive the **tumour**

HAIR FOLLICLE

A small sac in the skin from which hair grows from

HEPATIC

Relating to the liver

HISTOLOGICAL SUBTYPE

Cancer type based on the type of tissue in which the cancer started

HYPERCALCAEMIA

An abnormally high level of calcium in the blood

HYPERKALAEMIA

An abnormally high level of potassium in the blood

HYPERTENSION

Abnormally high blood pressure

HYPOCALCAEMIA

An abnormally low level of calcium in the blood

HYPOKALAEMIA

An abnormally low level of potassium in the blood

HYPOMAGNEAEMIA

An abnormally low level of magnesium in the blood

HYPONATRAEMIA

An abnormally low level of sodium in the blood

HYPOPHOSPHATAEMIA

An abnormally low level of phosphate in the blood

IMMUNOTHERAPY

A type of cancer treatment that stimulates the body's immune system to fight the cancer

INDUCTION THERAPY

Initial treatment with **chemotherapy** and/or **radiotherapy** to shrink the **tumour** before a second planned treatment (for example, surgery)

INTRAVENOUS

Administered into a vein

IONISING RADIATION

Any type of particle or electromagnetic wave that carries enough energy to ionize or remove electrons from an atom (e.g. **x-rays**)

LARGE CELL (UNDIFFERENTIATED) CARCINOMA

A type of NSCLC that does not look like **adenocarcinoma** or **squamous cell carcinoma** under the microscope

LEUKOPENIA

A decrease in the number of leukocytes (a type of white blood cell) in the blood, which places individuals at increased risk of infection

LIQUID BIOPSY

Tests performed in blood samples or other body fluids to detect the presence of substances that have originated in a **tumour**, and therefore, indicate the presence of a cancer

GLOSSARY

LOBECTOMY

A type of lung cancer surgery in which one **lobe** of a lung is removed (The right lung has three lobes, and the left lung has two lobes)

LOBE

A (usually rounded) part of an organ that appears to be separate in some way from the rest of that organ

LOCAL ANAESTHETIC

A medication that causes reversible absence of pain sensation around the site of administration

LOCALLY ADVANCED

Cancer that has spread from where it started to nearby tissue or **lymph nodes**

LYMPH

The fluid that circulates throughout the **lymphatic system**; it contains infection-fighting white blood cells

LYMPH NODES

Small structures throughout the **lymphatic system** that work as filters for harmful substances, such as cancer cells or bacteria

LYMPHATIC SYSTEM

A network of tissues and organs that help rid the body of toxins, waste and other unwanted materials. The primary function of the lymphatic system is to transport **lymph**, a fluid containing infection-fighting white blood cells, throughout the body

LYMPHOPENIA

An abnormally low level of lymphocytes (a type of white blood cell) in the blood, which places individuals at increased risk of infection

MAGNETIC RESONANCE IMAGING (MRI)

A type of scan that uses strong magnetic fields and radio waves to produce detailed images of the inside of the body

MAINTENANCE TREATMENT

Treatment given after the initial cycles of **chemotherapy** with the aim of keeping the cancer under control

METACHRONOUS OLIGOMETASTASES

Oligometastases that appear following treatment for a **primary tumour**

METASTATIC

A cancer that has spread from its (primary) site of origin to different parts of the body

METASTASIS (METASTASES)

A cancerous **tumour** or growth that has originated from a **primary tumour**/growth in another part of the body (plural = metastases)

MONOCLONAL ANTIBODY

A type of **targeted therapy**. Monoclonal antibodies recognise and attach to specific proteins produced by cells. Each monoclonal antibody recognises one particular protein. They work in different ways depending on the protein they are targeting

MUCOSITIS

Inflammation and ulceration of the membranes lining the **gastrointestinal system**

MULTIDISCIPLINARY TEAM

A group of health care workers who are members of different disciplines (e.g. **oncologist**, **nurse specialist**, physiotherapist, **radiologist**) and provide specific services to the patient. The activities of the team are brought together using a care plan

MULTIMODAL THERAPY

A treatment approach that includes two or more treatment types – usually some combination of surgery, **chemotherapy** and **radiotherapy**

MUTATION

A permanent alteration in the DNA sequence that makes up a gene, such that the sequence differs from what is found in most people

MYALGIA

Pain in a muscle(s)

NECITUMUMAB

A **targeted therapy (monoclonal antibody)** that blocks activity of **epidermal growth factor receptor**. It is administered through a drip into a vein in your arm or chest and in combination with two types of **chemotherapy (gemcitabine and cisplatin)**

NEUROLOGICAL

Relating to the nerves and the nervous system

NEUTROPENIA

An abnormally low level of **neutrophils** in the blood, which increases risk of infection

NEUTROPHIL

A type of white blood cell that play an important role in fighting off infection

GLOSSARY

NINTEDANIB

A type of **targeted therapy** that blocks proteins called protein kinases, which are present in cancer cells and involved in cancer cell growth. It is administered as a twice-daily capsule

NIVOLUMAB

A type of **immunotherapy** that blocks a protein called PD-1 on the surface of certain immune cells called T-cells; this activates the T-cells to find and kill cancer cells. It is administered through a drip into a vein in your arm or chest

NURSE SPECIALIST

A nurse specialised in the care of patients with a certain condition (e.g. cancer)

OEDEMA

A build-up of fluid in the body which causes the affected tissue to become swollen

OESOPHAGITIS

Inflammation of the **oesophagus**

OESOPHAGUS

The food pipe; the tube that connects your throat with your stomach

OLIGOMETASTATIC DISEASE (OLIGOMETASTASES)

Cancer that has spread from its original site to a limited number of other sites/organs; disease progression may occur at these sites but without spread to additional organs (**oligometastases** can be described as either **synchronous** or **metachronous**)

ONCOLOGIST

A doctor who specialises in the medical management of cancer

OSIMERTINIB

A type of **targeted therapy** called a **tyrosine kinase inhibitor**, which works by blocking signals within cancer cells and stopping the action of **epidermal growth factor receptor**, causing cancer cells to die. It is administered as a once-daily tablet to patients who have previously been treated with another **tyrosine kinase inhibitor**

PACLITAXEL

A type of **chemotherapy** that is administered through a drip into a vein in your arm or chest

PALLIATIVE CARE

The care of patients with advanced, progressive illness. It focuses on providing relief from pain, symptoms and physical and emotional stress, without dealing with the cause of the condition

PASSIVE SMOKING

The inhalation of smoke by a person who is not actively smoking themselves

PATHOLOGIST

Doctor who diagnoses disease by examining cell and tissue samples

PEMBROLIZUMAB

A type of **immunotherapy** that blocks a protein called PD-1 on the surface of certain immune cells called T-cells; this activates the T-cells to find and kill cancer cells. It is administered through a drip into a vein in your arm or chest

PEMETREXED

A type of **chemotherapy** drug used to treat NSCLC, which is given **intravenously** (directly into your bloodstream through a vein in your arm or chest)

PERICARDIUM

The membrane that encloses the heart

PERIPHERAL NEUROPATHY

Damage to the nerves in the extremities of the body. Symptoms may include pain, sensitivity, numbness or weakness in the hands, feet or lower legs

PHARYNGITIS

Inflammation of the pharynx, which is in the back of the throat

PLATELETS

Tiny blood cells that help your body form clots to stop bleeding

PLATINUM-BASED

A class of **chemotherapy** that includes **cisplatin** and **carboplatin**

PLEURA

One of the two membranes around the lungs. These two membranes are called the visceral and parietal pleurae

PNEUMONECTOMY

The surgical removal of a lung or part of a lung

PNEUMONIA

Inflammation of the lung, usually caused by an infection

POSITRON EMISSION TOMOGRAPHY (PET)

An imaging test that uses a dye with radioactive tracers, which is injected into a vein in your arm

PRIMARY LUNG CANCER

A cancer that first started in the lungs

GLOSSARY

PRIMARY TUMOUR

The **tumour** where the cancer first started to grow

PROGNOSIS

The likely outcome of a medical condition

PROGRAMMED DEATH LIGAND-1 (PD-L1)

A cellular protein thought to be involved in helping the **tumour** to evade detection by the body's immune system

PRURITUS

Severe itching of the skin

RADIATION PNEUMONITIS

Symptoms of cough, fever and fullness of the chest that usually appear between 2 weeks and 6 months following **radiotherapy** but are usually temporary

RADIOACTIVE

A material that is unstable and spontaneously emits energy (radiation)

RADIOLOGICAL EXAMINATION

A test that uses **x-rays** or other medical imaging techniques to visualise the body and organs for the detection of signs of cancer or other abnormalities

RADIOLOGIST

A doctor specialised in diagnosing and treating disease and injury through the use of medical imaging techniques such as **x-rays**, **computed tomography**, **magnetic resonance imaging**, **positron emission tomography** and **ultrasound**

RADIOTHERAPY

Treatment involving the use of high-energy radiation, which is commonly used to treat cancer

RAMUCIRUMAB

A type of **targeted therapy** that blocks the action of **vascular endothelial growth factor**, and prevents the cancer cells from developing their own blood supply, thus helping to slow down **tumour** growth. It is administered through a drip into a vein in your arm or chest in combination with another type of **chemotherapy**

REGIONAL LYMPH NODES

Lymph nodes close to the **tumour**

RELAPSE

Return of a cancer or deterioration in a person's state of health

RENAL

Relating to the kidneys

RESECTABLE

Able to be removed (resected) by surgery

RHINITIS

Inflammation of the lining inside the nose

SECOND LINE

The second treatment given to a patient once the initial (**first-line**) therapy has not worked or has been stopped because of the occurrence of side effects or other concerns

SEGMENT (OR WEDGE) RESECTION

Surgical removal of the segment of the lung where the **tumour** is located

SEQUENTIALLY

Treatment given one after the other

SQUAMOUS CELL CARCINOMA (SCC)

A type of NSCLC; usually occurs in the central part of the lung or in one of the **bronchi**

STENT

A small tube that is used to keep an airway or artery open

STEREOTACTIC ABLATIVE RADIOTHERAPY (SABR)

A specialised type of **radiotherapy** that is given to the **tumour** from many different directions using detailed scans to ensure precise targeting so that higher doses can be given over a shorter time

STOMATITIS

Inflammation of the inside of the mouth

SUPPORTIVE CARE

Care that provides relief from pain, symptoms and physical and emotional stress, without treating the cancer itself

SYNCHRONOUS OLIGOMETASTASES

Oligometastases diagnosed within a few months of a **primary tumour**

SYSTEMIC ANTICANCER TREATMENT

Drugs that spread throughout the body to treat cancer cells wherever they may be. They include **chemotherapy**, hormonal therapy, **targeted therapy**, and **immunotherapy**

T790M MUTATION

A mutation of the **epidermal growth factor receptor** (also known as Threonine 790 Methionine [Thr790Met] mutation)

GLOSSARY

TARGETED THERAPY

A newer type of cancer treatment that uses drugs or other substances to precisely identify and attack cancer cells, usually while doing little damage to normal cells

TAXANE

A class of **chemotherapy** that includes **paclitaxel** and **docetaxel**

THIRD-LINE

A third line of treatment given to a patient once the previous two lines (**first-line** and **second-line**) of therapy have not worked or have been stopped because of the occurrence of side effects or other concerns

THROMBOCYTOPENIA

A deficiency of **platelets** in the blood. This causes bleeding into the tissues, bruising, and slow blood clotting after injury

THROMBOSIS

The formation of a blood clot inside a blood vessel, obstructing the flow of blood through the blood system

TINNITUS

The hearing of a sound (such as ringing, whining or buzzing) when no external sound is present

TRACHEA

The windpipe – the wide, hollow tube that connects the larynx (or voice box) to the **bronchi** of the lungs

TUMOUR

A lump or growth of abnormal cells. **Tumours** may be benign (not cancerous) or malignant (cancerous). In this guide, the term '**tumour**' refers to a cancerous growth, unless otherwise stated

TYROSINE KINASE INHIBITOR (TKI)

A type of **targeted therapy** that inhibits tyrosine kinases, which are substances that send growth signals to cells

ULTRASOUND

A type of medical scan where sound waves are converted into images by a computer

UNRESECTABLE

Unable to be removed (resected) by surgery

URANIUM

A naturally radioactive element

VASCULAR ENDOTHELIAL GROWTH FACTOR (VEGF)

A protein produced by cells that stimulates the growth of new **blood vessels**

VIDEO-ASSISTED THORACIC SURGERY (VATS)

A surgical procedure that allows doctors to see inside the chest and lungs. It is a form of 'keyhole' surgery

VINORELBINE

A type of **chemotherapy** that is administered through a drip into a vein in your arm or chest

WEDGE (OR SEGMENT) RESECTION

Surgical removal of the segment of the lung where the **tumour** is located

X-RAY

An imaging test, using a type of radiation that can pass through the body, that allows your doctor to see inside your body

Non-small-cell lung cancer

This guide has been prepared to help you, your friends and your family better understand the nature of non-small-cell lung cancer (NSCLC) and the treatments that are available. The medical information described in this document is based on the clinical practice guidelines of the European Society for Medical Oncology (ESMO) for the management of early-stage, locally advanced or metastatic NSCLC. We recommend that you ask your doctor about the tests and types of treatments available in your country for your type and stage of NSCLC.

This guide has been written by Kstorfin Medical Communications Ltd on behalf of ESMO.

© Copyright 2017 European Society for Medical Oncology. All rights reserved worldwide.

European Society for Medical Oncology (ESMO)

Via L. Taddei 4

6962 Viganello-Lugano

Switzerland

Tel: +41 (0)91 973 19 99

Fax: +41 (0)91 973 19 02

E-mail: clinicalguidelines@esmo.org

We can help you understand non-small-cell lung cancer and the available treatment options.

The ESMO Guides for Patients are designed to assist patients, their relatives and caregivers to understand the nature of different types of cancer and evaluate the best available treatment choices. The medical information described in the Guides for Patients is based on the ESMO Clinical Practice Guidelines, which are designed to guide medical oncologists in the diagnosis, follow-up and treatment in different cancer types.

For more information, please visit www.esmo.org

